

Before we begin, are there any questions from last day's work?
check the graphs from 2.3.3 and 2.5.3

Today's Learning Goal(s):

By the end of the class, I will be able to:

- a) verify the zeros of cubics and quartics are correct by graphing with technology
- b) describe and sketch the graph of a polynomial function from its key properties (i.e. zeros, end behaviour, the shape of the graph)
- c) expand the factored form of a function to verify it is the same as the function in standard form
- d) connect the zeros of the function with the x -intercepts of the graph

2.6.1: To Agree or Disagree...

Date: _____

Anticipation Guide


Instructions:

Check "Agree" or "Disagree" beside each statement *before* you start the task on BLM 2.6.2.

Compare your choice and explanation with a partner.

Revisit your choices after completing the task on BLM 2.6.2.

Compare the choices you made before the task and after the task.













Before		Statement	After	
Agree	Disagree		Agree	Disagree
2	11	1. The x-intercepts of the function $y = 2(x+1)(x-3)(x-4)$ are -1, 3, 4.		✓
15	1	2. If the end behaviour of a function is as $x \rightarrow \infty, y \rightarrow \infty$, the function has a positive leading coefficient.	✓	
5	9	3. The function $y = 2(x-1)(x+2)(x-3)$ has the same x-intercepts as $y = -5(x-1)(x+2)(x-3)$.	✓	
1	14	4. The function shown can be expressed in the form  $y = ax^4 + bx^3 + cx^2 + dx + e$		✓
8	6	5. There are an infinite number of cubic functions that have x-intercepts of -2, 0, and 3.	✓	
0	13	6. The zeros of the functions $y = x^4 - 5x^2 + 4$ and $y = (x+1)(x-1)(x+2)(x-2)$ are the same.	✓	

$$\begin{aligned}
 y &= x^4 - 5x^2 + 4 \\
 &= (x^2 - 4)(x^2 - 1) \\
 &= (x-2)(x+2)(x-1)(x+1)
 \end{aligned}$$

2.6.2: “Expanding” Your Understanding of Functions

Date: _____

Student Instructions: In your groups, match the graph of the six functions given in the envelope with the zeros, end behaviour, and defining equation in factored form.
Optional: Also match the defining equation in standard form.

Graph of Function	Zeros	End Behaviours	Defining Equation in Factored Form	Defining Equation in Standard Form (Optional)
	 -2, 0, 3	 as $x \rightarrow -\infty, y \rightarrow -\infty$ and as $x \rightarrow \infty, y \rightarrow \infty$	 $y = 5x(x - 3)(x + 2)$	$y = 5x^3 - 5x^2 - 30x$
	 -2, 0, 3	 as $x \rightarrow -\infty, y \rightarrow -\infty$ and as $x \rightarrow \infty, y \rightarrow \infty$	 $y = 2x(x - 3)(x + 2)$	$y = 2x^3 - 2x^2 - 12x$
	 -3, 0, 2	 as $x \rightarrow -\infty, y \rightarrow -\infty$ and as $x \rightarrow \infty, y \rightarrow \infty$	 $y = 2x(x + 3)(x - 2)$	$y = 2x^3 + 2x^2 - 12x$

2.6.1-2.6.3 Determining Equations of Polynomial Functions (Spring 2017)-s17 March 6, 2017

Graph of Function	Zeros	End Behaviours	Defining Equation in Factored Form	Defining Equation in Standard Form (Optional)
<pre> WINDOW Xmin=-8 Xmax=8 Xscl=1 Ymin=-100 Ymax=120 Yscl=20 Xres=1 </pre>	<p>-2, -1, 0, 3</p>	<p>as $x \rightarrow -\infty, y \rightarrow \infty$ and as $x \rightarrow \infty, y \rightarrow \infty$</p>	<p>$y = 2x(x+2)(x+1)(x-3)$</p>	<p>$y = 2x^4 - 14x^2 - 12x$</p>
<pre> WINDOW Xmin=-8 Xmax=8 Xscl=1 Ymin=-40 Ymax=120 Yscl=20 Xres=1 </pre>	<p>-3, -2, 0, 2</p>	<p>as $x \rightarrow -\infty, y \rightarrow \infty$ and as $x \rightarrow \infty, y \rightarrow \infty$</p>	<p>$y = 2x(x+3)(x+2)(x-2)$</p>	<p>$y = 2x^4 + 6x^3 - 8x^2 - 24x$</p>
<pre> WINDOW Xmin=-8 Xmax=8 Xscl=1 Ymin=-100 Ymax=120 Yscl=20 Xres=1 </pre>	<p>-3, 0, 2, 3</p>	<p>as $x \rightarrow -\infty, y \rightarrow \infty$ and as $x \rightarrow \infty, y \rightarrow \infty$</p>	<p>$y = 2x(x+3)(x-2)(x-3)$</p>	<p>$y = 2x^4 - 4x^3 - 18x^2 + 36x$</p>

Ex. 1 (on next page)

Determining Equations of Polynomial Functions (using the roots and a point)

Ex. 1 a) Write the equation of the “family” of quadratic functions with zeros -4 and 1 .

b) Determine the equation of the “member of the family” that passes through $(-3, 2)$.

Solution (You’ve done these types of questions (with quadratics) in previous grades)

a) The “family” of quadratic functions with zeros -4 and 1 is $y = a(x+4)(x-1)$.

All members of the same “family” will have the same zeros, but different values for a , and therefore different graphs (which are all vertical stretches or compressions of the graph if $a=1$). Read page 208 in the text for examples, and nice colour sketches.

b) To determine the equation of the **SPECIFIC** “member of the family” that passes through $(-3, 2)$, simply substitute the known value $(-3, 2)$ into the family equation, and solve for “ a ”.

$$y = a(x+4)(x-1)$$

$$(2) = a((-3)+4)((-3)-1)$$

$$2 = a(1)(-4)$$

$$2 = -4a$$

$$\frac{2}{-4} = a$$

$$\therefore a = -\frac{1}{2}$$

$$\text{and the equation is } y = \frac{-1}{2}(x+4)(x-1)$$

You will use the same method for polynomial functions of degree 3 and 4.

Now **Read pp.209-211 (Ex. 1-3)**

Then complete:

1) 2.6.3

2) pp.212-214 #8, 10, 14bcd, 16, 17(a-d)